

Chronic changes in the atrophied submandibular gland after long-term ligation of the main excretory duct in mice

学位名	博士(歯学)
学位授与機関	日本歯科大学
学位授与年度	2014
学位授与番号	32667甲第1111号
URL	http://id.nii.ac.jp/1102/00000733/

顎下腺主導管長期結紮マウスにおける萎縮唾液腺の経時的変化

村山和義

Chronic changes in the atrophied submandibular gland after long-term
ligation of the main excretory duct in mice

MURAYAMA Kazuyoshi

日本歯科大学大学院新潟生命歯学研究科生命歯学専攻
臨床科学系・顎口腔全身病学分野・顎口腔全身関連治療学
指導：田中 彰教授

The Nippon Dental University
Graduate School of Life Dentistry at Niigata,
Course of Clinical Science,
Field of Oral and Maxillofacial Surgery
Director: Prof. TANAKA Akira

2015 年 1 月

顎下腺主導管長期結紮マウスにおける萎縮唾液腺の経時的変化

村山和義

**Chronic changes in the atrophied submandibular gland after long-term ligation
of the main excretory duct in mice**

KAZUYOSHI Murayama

Abstract

To examine the pathology of salivary glands that have undergone atrophy or hypofunction due to old age or disease, the duct ligation model is used. This model has been used in studies on the course of glandular parenchyma atrophy and the potential for repair and regeneration. However, the period of ligation was short in most of these studies, and none have examined long-term progress. Therefore, we investigated long-term ligation of the submandibular gland in mice and examined the chronic changes in atrophied salivary glands.

The ligation periods were 1, 2 and 3 months, and atrophied salivary glands were resected after each period. Resected glands were examined by histology, immunohistochemistry, RT-PCR, and transmission electron microscopy. Histologically, disappearance of acinar cells and increases in duct-like structures were observed over time in atrophied salivary glands, resulting from long-term submandibular gland main excretory duct ligation. Acinar cell markers (α -amylase, aquaporin 5) showed marked weakness in expression after ligation, but expression was still observed after 3 months of ligation. Stem cell markers (c-kit, Sca-1) showed greater expression at 1 month of ligation, compared with controls, but expression subsequently decreased with time. Expression of the precursor cell marker cytokeratin 5 was retained throughout long-term ligation.

Atrophied salivary gland tissue resulting from long-term ligation showed increases in specific duct-like structures over time, and the expression of stem cell markers and progenitor cell markers in the area of these structures suggests that the repair capability remained intact.

Keywords: duct-like structure • duct ligation • c-kit • Sca-1 • cytokeratin 5